

# Functions

## CHAPTER 1

### Concepts and Skills

In this chapter you will review precalculus topics. Although these topics are not directly tested on the AP exam, reviewing them will reinforce some basic principles:

- general properties of functions: domain, range, composition, inverse;
- special functions: absolute value, greatest integer; polynomial, rational, trigonometric, exponential, and logarithmic;

and the BC topic,

- parametrically defined curves

one input to one output  
function

one input → two outputs x

Function is a **RELATIONSHIP** between two variables such that one variable is matched **UNIQUELY** with the other variable.

### A. DEFINITIONS

**A1.** A function  $f$  is a correspondence that associates with each element  $a$  of a set called the *domain* one and only one element  $b$  of a set called the *range*. We write

$$f(a) = b$$

to indicate that  $b$  is the *value* of  $f$  at  $a$ . The elements in the domain are called *inputs*, and those in the range are called *outputs*.

A function is often represented by an equation, a graph, or a table.

A vertical line cuts the graph of a function in at most one point.

Function  
Domain  
Range

x	y		
1	3	1	5
2	4	2	6
3	5	2	7
4	6	3	8
		4	9

### EXAMPLE 1

The **domain** of  $f(x) = x^2 - 2$  is the set of all real numbers; its **range** is the set of all reals greater than or equal to  $-2$ . Note that

$$f(0) = 0^2 - 2 = -2, \quad f(-1) = (-1)^2 - 2 = -1,$$

$$f(\sqrt{3}) = (\sqrt{3})^2 - 2 = 1, \quad f(c) = c^2 - 2,$$

$$f(x+h) - f(x) = [(x+h)^2 - 2] - [x^2 - 2]$$

$$= x^2 + 2hx + h^2 - 2 - x^2 + 2 = 2hx + h^2.$$

The set of values of  $x$

The set of values of  $y$

### EXAMPLE 2

Find the domains of: (a)  $f(x) = \frac{4}{x-1}$ ; (b)  $g(x) = \frac{x}{x^2-9}$ ; (c)  $h(x) = \frac{\sqrt{4-x}}{x}$ .

**SOLUTIONS:**

(a) The domain of  $f(x) = \frac{4}{x-1}$  is the set of all reals except  $x = 1$  (which we shorten to " $x \neq 1$ ").

(b) The domain of  $g(x) = \frac{x}{x^2-9}$  is  $x \neq 3, -3$ .

(c) The domain of  $h(x) = \frac{\sqrt{4-x}}{x}$  is  $x \leq 4, x \neq 0$  (which is a short way of writing  $\{x \mid x \text{ is real, } x < 0 \text{ or } 0 < x \leq 4\}$ ).



**A2.** Two functions  $f$  and  $g$  with the same domain may be combined to yield their sum and difference:  $f(x) + g(x)$  and  $f(x) - g(x)$ , also written as  $(f + g)(x)$  and  $(f - g)(x)$ , respectively; or their product and quotient:  $f(x)g(x)$  and  $f(x)/g(x)$ , also written as  $(fg)(x)$  and  $(f/g)(x)$ , respectively. The quotient is defined for all  $x$  in the shared domain except those values for which  $g(x)$ , the denominator, equals zero.

**EXAMPLE 3**

If  $f(x) = x^2 - 4x$  and  $g(x) = x + 1$ , then find  $\frac{f(x)}{g(x)}$  and  $\frac{g(x)}{f(x)}$ .

**SOLUTIONS:**  $\frac{f(x)}{g(x)} = \frac{x^2 - 4x}{x + 1}$  and has domain  $x \neq -1$ ;

$\frac{g(x)}{f(x)} = \frac{x + 1}{x^2 - 4x} = \frac{x + 1}{x(x - 4)}$  and has domain  $x \neq 0, 4$ .

**Composition**

**A3.** The *composition* (or *composite*) of  $f$  with  $g$ , written as  $f(g(x))$  and read as “ $f$  of  $g$  of  $x$ ,” is the function obtained by replacing  $x$  wherever it occurs in  $f(x)$  by  $g(x)$ . We also write  $(f \circ g)(x)$  for  $f(g(x))$ . The domain of  $(f \circ g)(x)$  is the set of all  $x$  in the domain of  $g$  for which  $g(x)$  is in the domain of  $f$ .

**EXAMPLE 4A**

If  $f(x) = 2x - 1$  and  $g(x) = x^2$ , then does  $f(g(x)) = g(f(x))$ ?

**SOLUTION:**  $f(g(x)) = 2(x^2) - 1 = 2x^2 - 1$

$g(f(x)) = (2x - 1)^2 = 4x^2 - 4x + 1$ .

In general,  $f(g(x)) \neq g(f(x))$ .

**EXAMPLE 4B**

If  $f(x) = 4x^2 - 1$  and  $g(x) = \sqrt{x}$ , find  $f(g(x))$  and  $g(f(x))$ . **composite**

**SOLUTIONS:**  $f(g(x)) = 4x - 1$  ( $x \geq 0$ );  $g(f(x)) = \sqrt{4x^2 - 1}$  ( $|x| \geq \frac{1}{2}$ ).

**Symmetry**

**A4.** A function  $f$  is **odd** if, for all  $x$  in the domain of  $f$ ,  $f(-x) = -f(x)$ .  
**even** if, for all  $x$  in the domain of  $f$ ,  $f(-x) = f(x)$ .

The graph of an odd function is symmetric about the origin; the graph of an even function is symmetric about the y-axis.



### EXAMPLE 5

The graphs of  $f(x) = \frac{1}{2}x^3$  and  $g(x) = 3x^2 - 1$  are shown in Figure N1-1;  $f(x)$  is odd,  $g(x)$  even.




FIGURE N1-1

**A5.** If a function  $f$  yields a single output for each input and also yields a single input for every output, then  $f$  is said to be *one-to-one*. Geometrically, this means that any horizontal line cuts the graph of  $f$  in at most one point. The function sketched at the left in Figure N1-1 is one-to-one; the function sketched at the right is not. A function that is increasing (or decreasing) on an interval  $I$  is one-to-one on that interval (see pages 162-163 for definitions of increasing and decreasing functions).

**A6.** If  $f$  is one-to-one with domain  $X$  and range  $Y$ , then there is a function  $f^{-1}$ , with domain  $Y$  and range  $X$ , such that

$$f^{-1}(y_0) = x_0 \quad \text{if and only if} \quad f(x_0) = y_0.$$

The function  $f^{-1}$  is the *inverse* of  $f$ . It can be shown that  $f^{-1}$  is also one-to-one and that its inverse is  $f$ . The graphs of a function and its inverse are symmetric with respect to the line  $y = x$ .


To find the inverse of  $y = f(x)$ ,  
interchange  $x$  and  $y$ ,  
then solve for  $y$ .

### EXAMPLE 6


Find the inverse of the one-to-one function  $f(x) = x^3 - 1$ .

**SOLUTION:** Interchange  $x$  and  $y$ :  $x = y^3 - 1$

Solve for  $y$ :  $y = \sqrt[3]{x+1} = f^{-1}(x)$ .







Note that the graphs of  $f$  and  $f^{-1}$  in Figure N1-2 are mirror images, with the line  $y = x$  as the mirror.

## Zeros

**A7.** The *zeros* of a function  $f$  are the values of  $x$  for which  $f(x) = 0$ ; they are the  $x$ -intercepts of the graph of  $y = f(x)$ .

### EXAMPLE 7


Find zeros of  $f(x) = x^4 - 2x^2$ .

**SOLUTION:** The zeros are the  $x$ 's for which  $x^4 - 2x^2 = 0$ . The function has three zeros, since  $x^4 - 2x^2 = x^2(x^2 - 2)$  equals zero if  $x = 0$ ,  $+\sqrt{2}$ , or  $-\sqrt{2}$ .

## B. SPECIAL FUNCTIONS

The *absolute-value* function  $f(x) = |x|$  and the *greatest-integer* function  $g(x) = [x]$  are sketched in Figure N1-3.


**Absolute  
value**



**Greatest  
integer**

$$f(x) = |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Absolute-value function



$g(x) = [x]$  is the  
greatest integer  
not greater than  $x$

Greatest-integer function

**FIGURE N1-3**



## EXAMPLE 8

A function  $f$  is defined on the interval  $[-2, 2]$  and has the graph shown in Figure N1-4.

- (a) Sketch the graph of  $y = |f(x)|$ .
- (b) Sketch the graph of  $y = f(|x|)$ .
- (c) Sketch the graph of  $y = -f(x)$ .
- (d) Sketch the graph of  $y = f(-x)$ .




FIGURE N1-4

**SOLUTIONS:** The graphs are shown in Figures N1-4a through N1-4d.




FIGURE N1-4a




FIGURE N1-4b






FIGURE N1-4c

FIGURE N1-4d

Note that graph (c) of  $y = -f(x)$  is the reflection of  $y = f(x)$  in the  $x$ -axis, whereas Figure graph (d) of  $y = f(-x)$  is the reflection of  $y = f(x)$  in the  $y$ -axis. How do the graphs of  $|f(x)|$  and  $f(|x|)$  compare with the graph of  $f(x)$ ?

## EXAMPLE 9

Let  $f(x) = x^3 - 3x^2 + 2$ . Graph the following functions on your calculator in the window  $[-3, 3] \times [-3, 3]$ : (a)  $y = f(x)$ ; (b)  $y = |f(x)|$ ; (c)  $y = f(|x|)$ .

**SOLUTIONS:**

(a)  $y = f(x)$

See Figure N1-5a.




FIGURE N1-5a

(b)  $y = |f(x)|$

See Figure N1-5b.




FIGURE N1-5b



(c)  $y = f(|x|)$   
See Figure N1-5c.




FIGURE N1-5c

Note how the graphs for (b) and (c) compare with the graph for (a).

## C. POLYNOMIAL AND OTHER RATIONAL FUNCTIONS

### C1. Polynomial Functions.

A *polynomial function* is of the form

$$f(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n,$$

where  $n$  is a positive integer or zero, and the  $a_k$ 's, the *coefficients*, are constants. If  $a_0 \neq 0$ , the degree of the polynomial is  $n$ .

A *linear function*,  $f(x) = mx + b$ , is of the first degree; its graph is a straight line with slope  $m$ , the constant rate of change of  $f(x)$  (or  $y$ ) with respect to  $x$ , and  $b$  is the line's  $y$ -intercept.

A *quadratic function*,  $f(x) = ax^2 + bx + c$ , has degree 2; its graph is a parabola that opens up if  $a > 0$ , down if  $a < 0$ , and whose axis is the line  $x = -\frac{b}{2a}$ .

A *cubic*,  $f(x) = a_0x^3 + a_1x^2 + a_2x + a_3$ , has degree 3; calculus enables us to sketch its graph easily; and so on. The domain of every polynomial is the set of all reals.

Polynomial

### C2. Rational Functions.

A *rational function* is of the form

$$f(x) = \frac{P(x)}{Q(x)},$$

where  $P(x)$  and  $Q(x)$  are polynomials. The domain of  $f$  is the set of all reals for which  $Q(x) \neq 0$ .

Rational and irrational

p/q recurring  
decimals

pie, e, sq rt of nonsqua

Rational  
function

## D. TRIGONOMETRIC FUNCTIONS

The fundamental trigonometric identities, graphs, and reduction formulas are given in the Appendix.



Trigonometric  
functions**D1. Periodicity and Amplitude.**

The trigonometric functions are periodic. A function  $f$  is *periodic* if there is a positive number  $p$  such that  $f(x + p) = f(x)$  for each  $x$  in the domain of  $f$ . The smallest such  $p$  is called the *period* of  $f$ . The graph of  $f$  repeats every  $p$  units along the  $x$ -axis. The functions  $\sin x$ ,  $\cos x$ ,  $\csc x$ , and  $\sec x$  have period  $2\pi$ ;  $\tan x$  and  $\cot x$  have period  $\pi$ .

The function  $f(x) = A \sin bx$  has amplitude  $A$  and period  $\frac{2\pi}{b}$ ;  $g(x) = \tan cx$  has period  $\frac{\pi}{c}$ .

**EXAMPLE 10**

Consider the function  $f(x) = \frac{1}{k} \cos(kx)$ .

- (a) For what value of  $k$  does  $f$  have period 2?  
 (b) What is the amplitude of  $f$  for this  $k$ ?

**SOLUTIONS:**

- (a) Function  $f$  has period  $\frac{2\pi}{k}$ ; since this must equal 2, we solve the equation

$$\frac{2\pi}{k} = 2, \text{ getting } k = \pi.$$

- (b) It follows that the amplitude of  $f$  that equals  $\frac{1}{k}$  has a value of  $\frac{1}{\pi}$ .

**EXAMPLE 11**

Consider the function  $f(x) = 3 - \sin \frac{\pi x}{3}$ .

Find (a) the period and (b) the maximum value of  $f$ .

- (c) What is the smallest positive  $x$  for which  $f$  is a maximum?  
 (d) Sketch the graph.


**SOLUTIONS:**

- (a) The period of  $f$  is  $2\pi \div \frac{\pi}{3}$ , or 6.

- (b) Since the maximum value of  $-\sin x$  is  $-(-1)$  or  $+1$ , the maximum value of  $f$  is  $3 + 1$  or 4.

- (c)  $-(\sin \frac{\pi x}{3})$  equals  $+1$  when  $\sin \frac{\pi x}{3} = -1$ , that is, when  $\frac{\pi x}{3} = \frac{3\pi}{2}$ . Solving yields  $x = \frac{9}{2}$ .

- (d) We graph  $y = 3 - \sin \frac{\pi x}{3}$  in  $[-5, 8] \times [0, 5]$ :

**FIGURE N1-6**



## D2. Inverses.

We obtain *inverses* of the trigonometric functions by limiting the domains of the latter so each trigonometric function is one-to-one over its restricted domain. For example, we restrict

$$\sin x \text{ to } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2},$$

$$\cos x \text{ to } 0 \leq x \leq \pi,$$

$$\tan x \text{ to } -\frac{\pi}{2} < x < \frac{\pi}{2}.$$

### Inverse trig functions

The graphs of  $f(x) = \sin x$  on  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$  and of its inverse  $f^{-1}(x) = \sin^{-1}x$  are shown in Figure N1-7. The inverse trigonometric function  $\sin^{-1}x$  is also commonly denoted by  $\arcsin x$ , which denotes *the* angle whose sine is  $x$ . The graph of  $\sin^{-1}x$  is, of course, the reflection of the graph of  $\sin x$  in the line  $y = x$ .




FIGURE N1-7

Also, for other inverse trigonometric functions,

$y = \cos^{-1}x$  (or  $\arccos x$ ) has domain  $-1 \leq x \leq 1$  and range  $0 \leq y \leq \pi$ ;

$y = \tan^{-1}x$  (or  $\arctan x$ ) has domain the set of reals and range  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .

Note also that

$$\sec^{-1}(x) = \cos^{-1}\left(\frac{1}{x}\right), \quad \csc^{-1}(x) = \sin^{-1}\left(\frac{1}{x}\right), \quad \text{and} \quad \cot^{-1}(x) = \frac{\pi}{2} - \tan^{-1}(x).$$



## E. EXPONENTIAL AND LOGARITHMIC FUNCTIONS

### E1. Exponential Functions.

#### Exponential functions

The following laws of exponents hold for all rational  $m$  and  $n$ , provided that  $a > 0$ ,  $a \neq 1$ :

$$\begin{aligned} a^0 &= 1; & a^1 &= a; & a^m \cdot a^n &= a^{m+n}; & a^m \div a^n &= a^{m-n}; \\ (a^m)^n &= a^{mn}; & a^{-m} &= \frac{1}{a^m}. \end{aligned}$$

The exponential function  $f(x) = a^x$  ( $a > 0$ ,  $a \neq 1$ ) is thus defined for all real  $x$ ; its domain is the set of positive reals. The graph of  $y = a^x$ , when  $a = 2$ , is shown in Figure N1-8.

Of special interest and importance in the calculus is the exponential function  $f(x) = e^x$ , where  $e$  is an irrational number whose decimal approximation to five decimal places is 2.71828. We define  $e$  on page 97.

### E2. Logarithmic Functions.

#### Log functions

Since  $f(x) = a^x$  is one-to-one, it has an inverse,  $f^{-1}(x) = \log_a x$ , called the *logarithmic function* with base  $a$ . We note that

$$y = \log_a x \quad \text{if and only if} \quad a^y = x.$$

The domain of  $\log_a x$  is the set of positive reals; its range is the set of all reals. It follows that the graphs of the pair of mutually inverse functions  $y = 2^x$  and  $y = \log_2 x$  are symmetric to the line  $y = x$ , as can be seen in Figure N1-8.




FIGURE N1-8

The logarithmic function  $\log_a x$  ( $a > 0$ ,  $a \neq 1$ ) has the following properties:

$$\begin{aligned} \log_a 1 &= 0; & \log_a a &= 1; & \log_a mn &= \log_a m + \log_a n; \\ \log_a \frac{m}{n} &= \log_a m - \log_a n; & \log_a x^m &= m \log_a x. \end{aligned}$$



The logarithmic base  $e$  is so important and convenient in calculus that we use a special symbol:

$$\log_e x = \ln x.$$

Logarithms with base  $e$  are called *natural* logarithms. The domain of  $\ln x$  is the set of positive reals; its range is the set of all reals. The graphs of the mutually inverse functions  $\ln x$  and  $e^x$  are given in the Appendix.

## F. PARAMETRICALLY DEFINED FUNCTIONS

If the  $x$ - and  $y$ -coordinates of a point on a graph are given as functions  $f$  and  $g$  of a third variable, say  $t$ , then

$$x = f(t), \quad y = g(t)$$

are called *parametric equations* and  $t$  is called the *parameter*. When  $t$  represents time, as it often does, then we can view the curve as that followed by a moving particle as the time varies.

BC ONLY

Parametric equations

### EXAMPLE 12

Find the Cartesian equation of, and sketch, the curve defined by the parametric equations

$$x = 4 \sin t, \quad y = 5 \cos t \quad (0 \leq t \leq 2\pi).$$

**SOLUTION:** We can eliminate the parameter  $t$  as follows:

$$\sin t = \frac{x}{4}, \quad \cos t = \frac{y}{5}.$$

Since  $\sin^2 t + \cos^2 t = 1$ , we have

$$\left(\frac{x}{4}\right)^2 + \left(\frac{y}{5}\right)^2 = 1 \quad \text{or} \quad \frac{x^2}{16} + \frac{y^2}{25} = 1$$

The curve is the ellipse shown in Figure N1-9.




FIGURE N1-9

Note that, as  $t$  increases from 0 to  $2\pi$ , a particle moving in accordance with the given parametric equations starts at point  $(0, 5)$  (when  $t = 0$ ) and travels in a clockwise direction along the ellipse, returning to  $(0, 5)$  when  $t = 2\pi$ .



## BC ONLY

## EXAMPLE 13

Given the pair of parametric equations,

$$x = 1 - t, \quad y = \sqrt{t} \quad (t \geq 0),$$

write an equation of the curve in terms of  $x$  and  $y$ , and sketch the graph.

**SOLUTION:** We can eliminate  $t$  by squaring the second equation and substituting for  $t$  in the first; then we have

$$y^2 = t \quad \text{and} \quad x = 1 - y^2.$$

We see the graph of the equation  $x = 1 - y^2$  on the left in Figure N1-10. At the right we see only the upper part of this graph, the part defined by the parametric equations for which  $t$  and  $y$  are both restricted to nonnegative numbers.




FIGURE N1-10

The function defined by the parametric equations here is  $y = F(x) = \sqrt{1 - x}$ , whose graph is at the right above; its domain is  $x \leq 1$  and its range is the set of nonnegative reals.

## EXAMPLE 14

A satellite is in orbit around a planet that is orbiting around a star. The satellite makes 12 orbits each year. Graph its path given by the parametric equations

$$x = 4 \cos t + \cos 12t,$$

$$y = 4 \sin t + \sin 12t.$$

**SOLUTION:** Shown below is the graph of the satellite's path using the calculator's parametric mode for  $0 \leq t \leq 2\pi$ .




FIGURE N1-11



**EXAMPLE 15**

Graph  $x = y^2 - 6y + 8$ .

**SOLUTION:** We encounter a difficulty here. The calculator is constructed to graph  $y$  as a function of  $x$ : it accomplishes this by scanning horizontally across the window and plotting points in varying vertical positions. Ideally, we want the calculator to scan *down* the window and plot points at appropriate horizontal positions. But it won't do that.

One alternative is to interchange variables, entering  $x$  as  $Y_1$  and  $y$  as  $X$ , thus entering  $Y_1 = X^2 - 6X + 8$ . But then, during all subsequent processing we must remember that we have made this interchange.

Less risky and more satisfying is to switch to parametric mode: Enter  $x = t^2 - 6t + 8$  and  $y = t$ . Then graph these equations in  $[-10, 10] \times [-10, 10]$ , for  $t$  in  $[-10, 10]$ . See Figure N1-12.




FIGURE N1-12

**EXAMPLE 16**

Let  $f(x) = x^3 + x$ ; graph  $f^{-1}(x)$ .

**SOLUTION:** Recalling that  $f^{-1}$  interchanges  $x$  and  $y$ , we use parametric mode to graph

$$\begin{aligned} f: x = t, y = t^3 + t \\ \text{and } f^{-1}: x = t^3 + t, y = t. \end{aligned}$$

Figure N1-13 shows both  $f(x)$  and  $f^{-1}(x)$ .




FIGURE N1-13

Parametric equations give rise to vector functions, which will be discussed in connection with motion along a curve in Chapter 4.



## Chapter Summary

This chapter has reviewed some important precalculus topics. These topics are not directly tested on the AP exam; rather, they represent basic principles important in calculus. These include finding the domain, range and inverse of a function; and understanding the properties of polynomial and rational functions, trigonometric and inverse trig functions, and exponential and logarithmic functions.

For BC students, this chapter also reviewed parametrically defined functions.

## Practice Exercises

**Directions:** Answer these questions without using your calculator.

1. If  $f(x) = x^3 - 2x - 1$ , then  $f(-2) =$

- (A) -17 (B) -13 (C) -5 (D) -1 (E) 3

2. The domain of  $f(x) = \frac{x-1}{x^2+1}$  is

- (A) all  $x \neq 1$  (B) all  $x \neq 1, -1$  (C) all  $x \neq -1$   
 (D)  $x \geq 1$  (E) all reals

3. The domain of  $g(x) = \frac{\sqrt{x-2}}{x^2-x}$  is

- (A) all  $x \neq 0, 1$  (B)  $x \leq 2, x \neq 0, 1$  (C)  $x \leq 2$   
 (D)  $x \geq 2$  (E)  $x > 2$

4. If  $f(x) = x^3 - 3x^2 - 2x + 5$  and  $g(x) = 2$ , then  $g(f(x)) =$

- (A)  $2x^3 - 6x^2 - 2x + 10$  (B)  $2x^2 - 6x + 1$  (C) -6  
 (D) -3 (E) 2

5. With the functions and choices as in Question 4, which choice is correct for  $f(g(x))$ ?

HW 6. If  $f(x) = x^3 + Ax^2 + Bx - 3$  and if  $f(1) = 4$  and  $f(-1) = -6$ , what is the value of  $2A + B$ ?

- (A) 12 (B) 8 (C) 0 (D) -2  
 (E) It cannot be determined from the given information.

7. Which of the following equations has a graph that is symmetric with respect to the origin?

- (A)  $y = \frac{x-1}{x}$  (B)  $y = 2x^4 + 1$  (C)  $y = x^3 + 2x$   
 (D)  $y = x^3 + 2$  (E)  $y = \frac{x}{x^3+1}$

8. Let  $g$  be a function defined for all reals. Which of the following conditions is not sufficient to guarantee that  $g$  has an inverse function?

- (A)  $g(x) = ax + b, a \neq 0$ . (B)  $g$  is strictly decreasing.  
 (C)  $g$  is symmetric to the origin. (D)  $g$  is strictly increasing.  
 (E)  $g$  is one-to-one.

0 and

asymptote at  $x=0$

$-1/x$

$1-1/x$

shift up

3 turns

2 turns

$f(-x) = -f(x)$  odd  
 $f(-x) = f(x)$  even

symmetric to y-axis



9. Let  $y = f(x) = \sin(\arctan x)$ . Then the range of  $f$  is

- (A)  $\{y \mid 0 < y \leq 1\}$  (B)  $\{y \mid -1 < y < 1\}$  (C)  $\{y \mid -1 \leq y \leq 1\}$   
 (D)  $\left\{y \mid -\frac{\pi}{2} < y < \frac{\pi}{2}\right\}$  (E)  $\left\{y \mid -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}\right\}$

10. Let  $g(x) = |\cos x - 1|$ . The maximum value attained by  $g$  on the closed interval  $[0, 2\pi]$  is for  $x$  equal to

- (A)  $-1$  (B)  $0$  (C)  $\frac{\pi}{2}$  (D)  $2$  (E)  $\pi$

11. Which of the following functions is not odd?

- (A)  $f(x) = \sin x$  (B)  $f(x) = \sin 2x$  (C)  $f(x) = x^3 + 1$  (D)  $f(x) = \frac{x}{x^2 + 1}$  (E)  $f(x) = \sqrt[3]{2x}$   
 $f(-x) = -f(x)$  odd  $f(-x) = f(x)$  even  
 $f(-x) = -x^3 + 1 = -(x^3 - 1)$  not odd

12. The roots of the equation  $f(x) = 0$  are 1 and  $-2$ . The roots of  $f(2x) = 0$  are

- (A) 1 and  $-2$  (B)  $\frac{1}{2}$  and  $-1$  (C)  $-\frac{1}{2}$  and 1  
 (D) 2 and  $-4$  (E)  $-2$  and 4

13. The set of zeros of  $f(x) = x^3 + 4x^2 + 4x$  is

- (A)  $\{-2\}$  (B)  $\{0, -2\}$  (C)  $\{0, 2\}$  (D)  $\{2\}$  (E)  $\{2, -2\}$

14. The values of  $x$  for which the graphs of  $y = x + 2$  and  $y^2 = 4x$  intersect are

- (A)  $-2$  and 2 (B)  $-2$  (C) 2 (D) 0 (E) none of these

15. The function whose graph is a reflection in the  $y$ -axis of the graph of  $f(x) = 1 - 3^x$  is

- (A)  $g(x) = 1 - 3^{-x}$  (B)  $g(x) = 1 + 3^x$  (C)  $g(x) = 3^x - 1$   
 (D)  $g(x) = \log_3(x - 1)$  (E)  $g(x) = \log_3(1 - x)$

16. Let  $f(x)$  have an inverse function  $g(x)$ . Then  $f(g(x)) =$

- (A) 1 (B)  $x$  (C)  $\frac{1}{x}$  (D)  $f(x) \cdot g(x)$  (E) none of these

17. The function  $f(x) = 2x^3 + x - 5$  has exactly one real zero. It is between

- (A)  $-2$  and  $-1$  (B)  $-1$  and 0 (C) 0 and 1  
 (D) 1 and 2 (E) 2 and 3



18. The period of  $f(x) = \sin \frac{2\pi}{3} x$  is
- (A)  $\frac{1}{3}$  (B)  $\frac{2}{3}$  (C)  $\frac{3}{2}$  (D) 3 (E) 6
19. The range of  $y = f(x) = \ln(\cos x)$  is
- (A)  $\{y \mid -\infty < y \leq 0\}$  (B)  $\{y \mid 0 < y \leq 1\}$  (C)  $\{y \mid -1 < y < 1\}$   
(D)  $\left\{y \mid -\frac{\pi}{2} < y < \frac{\pi}{2}\right\}$  (E)  $\{y \mid 0 \leq y \leq 1\}$
20. If  $\log_b(3^b) = \frac{b}{2}$ , then  $b =$
- (A)  $\frac{1}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{1}{2}$  (D) 3 (E) 9
21. Let  $f^{-1}$  be the inverse function of  $f(x) = x^3 + 2$ . Then  $f^{-1}(x) =$
- (A)  $\frac{1}{x^3 - 2}$  (B)  $(x + 2)^3$  (C)  $(x - 2)^3$   
(D)  $\sqrt[3]{x + 2}$  (E)  $\sqrt[3]{x - 2}$
22. The set of  $x$ -intercepts of the graph of  $f(x) = x^3 - 2x^2 - x + 2$  is
- (A)  $\{1\}$  (B)  $\{-1, 1\}$  (C)  $\{1, 2\}$   
(D)  $\{-1, 1, 2\}$  (E)  $\{-1, -2, 2\}$
23. If the domain of  $f$  is restricted to the open interval  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ , then the range of  $f(x) = e^{\tan x}$  is
- (A) the set of all reals (B) the set of positive reals  
(C) the set of nonnegative reals (D)  $\{y \mid 0 < y \leq 1\}$   
(E) none of these
24. Which of the following is a reflection of the graph of  $y = f(x)$  in the  $x$ -axis?
- (A)  $y = -f(x)$  (B)  $y = f(-x)$  (C)  $y = |f(x)|$   
(D)  $y = f(|x|)$  (E)  $y = -f(-x)$
25. The smallest positive  $x$  for which the function  $f(x) = \sin\left(\frac{x}{3}\right) - 1$  is a maximum is
- (A)  $\frac{\pi}{2}$  (B)  $\pi$  (C)  $\frac{3\pi}{2}$  (D)  $3\pi$  (E)  $6\pi$



26.  $\tan \left( \arccos \left( -\frac{\sqrt{2}}{2} \right) \right) =$   
 (A)  $-1$  (B)  $-\frac{\sqrt{3}}{3}$  (C)  $-\frac{1}{2}$  (D)  $\frac{\sqrt{3}}{3}$  (E)  $1$
27. If  $f^{-1}(x)$  is the inverse of  $f(x) = 2e^{-x}$ , then  $f^{-1}(x) =$   
 (A)  $\ln \left( \frac{2}{x} \right)$  (B)  $\ln \left( \frac{x}{2} \right)$  (C)  $\left( \frac{1}{2} \right) \ln x$   
 (D)  $\sqrt{\ln x}$  (E)  $\ln (2 - x)$
28. Which of the following functions does not have an inverse function?  
 (A)  $y = \sin x \left( -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \right)$  (B)  $y = x^3 + 2$  (C)  $y = \frac{x}{x^2 + 1}$   
 (D)  $y = \frac{1}{2}e^x$  (E)  $y = \ln (x - 2)$  (where  $x > 2$ )
29. Suppose that  $f(x) = \ln x$  for all positive  $x$  and  $g(x) = 9 - x^2$  for all real  $x$ . The domain of  $f(g(x))$  is  
 (A)  $\{x \mid x \leq 3\}$  (B)  $\{x \mid |x| \leq 3\}$  (C)  $\{x \mid |x| > 3\}$   
 (D)  $\{x \mid |x| < 3\}$  (E)  $\{x \mid 0 < x < 3\}$
30. Suppose (as in Question 29) that  $f(x) = \ln x$  for all positive  $x$  and  $g(x) = 9 - x^2$  for all real  $x$ . The range of  $y = f(g(x))$  is  
 (A)  $\{y \mid y > 0\}$  (B)  $\{y \mid 0 < y \leq \ln 9\}$  (C)  $\{y \mid y \leq \ln 9\}$   
 (D)  $\{y \mid y < 0\}$  (E) none of these
31. The curve defined parametrically by  $x(t) = t^2 + 3$  and  $y(t) = t^2 + 4$  is part of a(n)  
 (A) line (B) circle (C) parabola  
 (D) ellipse (E) hyperbola
32. Which equation includes the curve defined parametrically by  $x(t) = \cos^2(t)$  and  $y(t) = 2 \sin(t)$ ?  
 (A)  $x^2 + y^2 = 4$  (B)  $x^2 + y^2 = 1$  (C)  $4x^2 + y^2 = 4$   
 (D)  $4x + y^2 = 4$  (E)  $x + 4y^2 = 1$

BC ONLY